

Sumagic®

PLANT GROWTH REGULATOR
COMMERCIAL
FOR GREENHOUSE USE ONLY

GUARANTEE: UNICONAZOLE-P.....0.055%

Contains 1,2-benzisothrazolin-3-one at 0.019% as a preservative

READ THE LABEL BEFORE USING

REGISTRATION NUMBER: 25781
PEST CONTROL PRODUCTS ACT

PLANT PRODUCTS CO. LTD.
314 ORENDA ROAD
BRAMPTON, ONTARIO L6T 1G1
800-898-2536

VALENT U.S.A CORPORATION
P.O. BOX 8025
Walnut Creek, CA 94596-8025
www.valentpro.com

CAUTION
POISON

NET CONTENTS 4 LITRES

PRECAUTIONS

CAUTION

- KEEP OUT OF REACH OF CHILDREN
- Avoid contact with eyes, skin and clothing.
- Wash concentrate from skin or eyes immediately.
- Do not inhale fumes. Avoid breathing spray mist. Avoid spray drift.
- After use, wash hands and other exposed skin.
- Remove and wash contaminated clothing before reuse.
- When mixing, loading and applying the product, wear waterproof rain gear (e.g. Tyvek coveralls) with a hood, respirator with an organic chemical cartridge, chemical resistant gloves, goggles and rubber boots.
- Do not reenter treated area within 12 hours of application. If early reentry is required, then wear waterproof rain gear (e.g. Tyvek coveralls), respirator with an organic chemical cartridge, chemical resistant gloves and rubber boots.
- Avoid contamination of ponds, streams, rivers and desirable vegetation.

- Store the container tightly closed and away from seed, feeds, fertilizers, plants and foodstuffs.

FIRST AID

If swallowed: If swallowed give water or milk to drink and call a physician or contact a poison control centre. Consult medical personnel before inducing vomiting. If medical advice cannot be obtained, then take the person and product container to the nearest medical emergency treatment centre or hospital. If person is unconscious do NOT give anything by mouth and do NOT induce vomiting.

If inhaled: If inhaled, remove victim to fresh air. If not breathing, give artificial respiration preferably mouth-to-mouth. Get medical attention or contact a poison control centre.

If in eyes: Flush eyes with plenty of water. Call a physician or a poison control centre if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention or contact a poison control centre if irritation persists.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

Note to Physicians: Emergency Information – call 1-800-892-0099.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

DIRECTIONS FOR USE

IMPORTANT NOTES

- SUMAGIC Plant Growth Regulator is an extremely active plant growth regulator. Extreme caution must be exercised in measuring, diluting and applying SUMAGIC Plant Growth Regulator. Excessive doses will result in severe growth retardation and possible adverse effects on flowering, plant appearance and marketability. Avoid spray drift onto non-target species.
- Apply to well-watered, vigorously growing plants. Do not mix with pesticides, fertilizers, or adjuvants.

PRECAUTIONS

- FOR GREENHOUSE USE ONLY.
- Do not apply SUMAGIC Plant Growth Regulator to any food crops.
- Do not use in greenhouses with soil floors consisting of sand or sandy loam soil.
- Do not reuse soil from plants treated with SUMAGIC Plant Growth Regulator.
- Do **NOT** use on cranesbill geranium (*Geranium spp*).
- It is recommended that SUMAGIC Plant Growth Regulator be applied during morning or late afternoon hours.
- SUMAGIC Plant Growth Regulator retards elongation almost immediately after application. Assessment of rate response can often be made about one week after treatment.

- For Poinsettias and Chrysanthemums only, if response to SUMAGIC Plant Growth Regulator is deemed to be inadequate following an initial application within the recommended rate range, a repeat application can be made.
- Differences in plant response to SUMAGIC Plant Growth Regulator due to plant surface area in relation to soil surface area are possible.
- Extreme temperatures can influence plant response to SUMAGIC Plant Growth Regulator. Within the recommended rate range, a higher rate may be required at higher temperatures than at low temperatures for the same effect.

GENERAL INFORMATION

SUMAGIC Plant Growth Regulator can be used to retard the growth of Poinsettias, Chrysanthemums, Easter Lilies and Geranium and Bedding Plant Species. Dilute with water according to the directions below. Make foliar applications with standard spray equipment. Effective dose is strongly dependent on the application volume. Rates expressed are based on a spray volume of 2 litres applied uniformly to 10 square metres of bench area. Low pressure and larger pressure droplets are recommended for best results.

Plant response to SUMAGIC Plant Growth Regulator is strongly influenced by cultural and environmental variables, such as growing media, water/fertilizer management, temperature, light, relative day and night temperatures, growing house composition, and other cultural practices. First time users of SUMAGIC Plant Growth Regulator should treat a limited number of plants at the lowest recommended rate. Rates may be adjusted within the recommended rate range as necessary. In particular, temperature can have a significant effect on the response from any given rate. Within the recommended rate range, a higher rate may be required at higher temperatures than at lower temperatures for the same effect.

SUMAGIC Plant Growth Regulator has been tested on a limited range of cultivars. Response to SUMAGIC Plant Growth Regulator has been determined to be influenced by genetic background. Growers should limit first use of SUMAGIC Plant Growth Regulator to a limited number of plants of each cultivar, prior to adoption as a cultural practice under their particular growing conditions.

POINSETTIA

Apply SUMAGIC Plant Growth Regulator as a foliar spray when the breaks are 3.5 to 5.0 cm in length (about 10 to 14 days after pinching). Use a rate of 4 to 16 mL/L (2 to 8 ppm) and apply uniformly at a spray volume of 2 litres per 10 square metres of bench area. Up to two repeat applications can be made at 7 day intervals until the desired growth characteristics are achieved. Bract size and diameter may be slightly reduced with treatment, and days to flowering may be delayed. Late season application (after short-day initiation) may have negative effects on bract size, plant quality, and marketability. **DO NOT APPLY TO POINSETTIAS AFTER THE INITIATION OF SHORT DAYS.**

CHRYSANTHEMUMS

Apply SUMAGIC Plant Growth Regulator as a foliar spray 7 to 14 days after pinching when the length of the breaks is 3.8 to 5.0 cm. Use a rate of 10 to 20 mL/L (5 to 10 ppm) and apply uniformly in a spray volume of 2 litres per 10 square metres of bench area. A

single repeat spray can be applied 7 to 21 days after the initial application to increase the level of growth reduction. Two applications may have more effect than the same amount of product applied as a single treatment. Alternatively, a solution of 40 mL/L (20 ppm) can be applied in a spray volume of 2 litres per 10 square metres of bench area, but DO NOT make a repeat application following the 20 ppm spray.

EASTER LILIES

Apply SUMAGIC Plant Growth Regulator as a foliar spray when shoots are 7.5 cm tall. Use a rate of 6 to 60 mL/L (3 to 30 ppm) and apply uniformly in a volume of 2 litres per 10 square metres of bench area. A single repeat spray of 3 to 5 ppm may be applied 2 to 3 weeks after initial spray of 3 to 5 ppm. DO NOT make a repeat application following a spray of greater than 5 ppm. Avoid late-season application. Insufficient height reduction may result following application of SUMAGIC Plant Growth Regulator to “Nellie White” Easter lilies grown under zero or negative DIF (day temperatures the same as or lower than night temperatures).

NOTE: Treatment with SUMAGIC Plant Growth Regulator may reduce the water requirement for Easter Lilies.

GERANIUM AND BEDDING PLANT SPECIES

Apply as a foliar spray.

Geranium and Bedding Plant Species for which use of SUMAGIC Plant Growth Regulator is Acceptable, with Stated Restrictions.

HOST SPECIES	ACCEPTABLE RATE (PPM UNICONAZOLE-P)*	ACCEPTABLE MAXIMUM NUMBER OF APPLICATIONS**	ACCEPTABLE APPLICATION TIMING***
BEDDING PLANTS			
Celosia (<i>Celosia plumosa</i>)	2.5 to 10	Two	5 to 10 cm height
Coleus (<i>Coleus sp.</i>)	5 to 10	One	2 to 5 cm height
Impatiens (<i>Impatiens walleriana</i> or <i>I. balsamina</i>)	2.5 to 10	Two	3 to 10 cm height
Marigold (<i>Calendula officinalis</i> , <i>Tagetes erectus</i> , <i>T. patula</i> , and <i>T. signata</i>)	5 to 15	Two	5 to 10 cm height
Pansy (<i>Viola wittrockiana</i>)	1 to 6	One; if > 3 ppm was applied Two; both applications of 1 to 3 ppm	3 to 5 cm height
Periwinkle (<i>Vinca minor</i>)	1 to 3	Two	5 to 10 cm height
Petunia (<i>Petunia hybrida</i>)	10 to 30	One	2 to 5 cm height
Salvia (<i>Salvia splendens</i> , <i>S. horminum</i> , <i>S. farinacea</i> , <i>S. coccinea</i> , and <i>S. sclarea</i>)	1 to 10	Two	3 to 10 cm height
Snapdragon (<i>Antirrhinum majus</i>)	10 to 30	One; if >15 ppm was applied Two; both applications of 10 to 15 ppm	5 to 10 cm height
Viola (<i>V. tricolor</i> , <i>V. cornuta</i> , and <i>V. williamsii</i>)	1 to 5	One	2 to 5 cm height
Geranium (<i>Zonal: Pelargonium hortorum</i> , and <i>Ivy: Pelargonium peltatum</i>)	2 to 4 if grown from seed; 2 to 8 if vegetatively propagated	Two	Use 2 to 4 ppm on seed geraniums when plant height is 5 to 16 cm. For vegetatively propagated geranium, apply 2 to 8 ppm to cuttings once rooting is established.

* SUMAGIC Plant Growth Regulator is extremely active at very low rates. Growers should be careful to initially use lower rates for younger seedlings. Application may be made by foliar spray only.

- ** To some crops (as indicated), a second application may be made as required, but no earlier than 7 days after the first application.
- *** The application timing proposed was 5 to 16 cm height for all proposed bedding plant species. The application timing shown in this column was supported by the submitted data, however, the applicant and registrant may further restrict the application timing, but only within the accepted application window.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE

Keep pesticide in original container. Do not put concentrate or dilute solutions into food or drink container. Store in cool, dry, and well ventilated place away from seed, feed, fertilizer or other pesticides. Do not store diluted spray.

DISPOSAL

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

SUMAGIC is a registered trademark of Valent U.S.A. Corp.
Plant Products Co. Ltd. is an authorized user of the trademark.

Form 1327-F

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.